

Standard noncommuting and commuting dilations of commuting tuples

B. V. RAJARAMA BHAT, TIRTHANKAR BHATTACHARYYA, AND SANTANU DEY

April 5, 2002

Abstract

We introduce a notion called ‘maximal commuting piece’ for tuples of Hilbert space operators. Given a commuting tuple of operators forming a row contraction there are two commonly used dilations in multivariable operator theory. Firstly there is the minimal isometric dilation consisting of isometries with orthogonal ranges and hence it is a noncommuting tuple. There is also a commuting dilation related with a standard commuting tuple on Boson Fock space. We show that this commuting dilation is the maximal commuting piece of the minimal isometric dilation. We use this result to classify all representations of Cuntz algebra $\mathcal{O}_n$ coming from dilations of commuting tuples.

KEY WORDS: Dilation, Commuting Tuples, Complete Positivity, Cuntz Algebra
MATHEMATICS SUBJECT CLASSIFICATION: 47A20, 47A13, 46L05, 47D25

1. INTRODUCTION

It is a well-known result due to Sz. Nagy that every contraction on a Hilbert space dilates to an isometry. There is a very natural generalization of this result to a class of operator tuples defined as follows.

DEFINITION 1. A *contractive n -tuple*, or a *row contraction* is a n -tuple $\underline{T} = (T_1, \dots, T_n)$ of bounded operators on a Hilbert space $\mathcal{H}$ such that $T_1 T_1^* + \dots + T_n T_n^* \leq I$.

Such tuples are known as row contractions as the condition is equivalent to having the operator $(T_1, \dots, T_n)$ from $\mathcal{H} \oplus \dots \oplus \mathcal{H}$ (n -times) to $\mathcal{H}$, as a contraction. It is possible to dilate contractive tuples to tuples of isometries with orthogonal ranges. Moreover, such a dilation is unique up to unitary equivalence, under a natural minimality condition just as in the one variable case. This dilation which we call as *the minimal isometric dilation* or as *standard noncommuting dilation* has been explored by many authors. Some ideas along this direction can already be seen in the early paper [13] of Davis. In more concrete form this dilation can be seen in the papers of Bunce [11] and Frazho [16, 17]. A real extensive study of this notion has been carried out by Popescu in a series of papers ([20]-[24], [2, 3]) and he has neat generalizations of many results from one variable situation.

Now suppose the tuple under consideration is a commuting tuple in the sense that $T_i T_j = T_j T_i$ for all $1 \leq i, j \leq n$. Then it is natural to wish for a dilation consisting of mutually commuting isometries. Unfortunately, such a dilation does not exist in general for $n \geq 3$ [19]. However, there is a dilation of commuting contractive tuples studied recently by Arveson [5]. In fact, this particular dilation was looked at by Drury [15] in his study of von Neumann inequality for tuples and similar ideas have been explored by Agler [1], Athavale [7] and others for different classes of operators using various reproducing kernels. We call this dilation of Drury and Arveson as *standard commuting dilation* of contractive commuting tuples. This dilation consists of a commuting tuple, but the constituent operators are not isometries. Then a natural question arises as to in what sense this dilation is canonical. Here in Section 3 as our main result, we show that the standard commuting dilation is the ‘maximal commuting tuple contained’ in the minimal isometric dilation. To begin with we make these words inside inverted commas precise by defining what we call as ‘maximal commuting piece’ of a tuple of operators. We can explore how the standard commuting dilation of the maximal commuting piece sits inside the minimal isometric dilation of the original tuple, and whether it is the maximal commuting piece of the minimal isometric dilation etc. We have been able to carry out this study for purely contractive tuples in Section 2.

Any tuple $(W_1, \dots, W_n)$ of isometries with orthogonal ranges satisfying $\sum W_i W_i^* = I$ gives us a concrete representation of the familiar Cuntz algebra [12]. Recently, there has been a lot of effort to study such representations in connection with wavelet theory, see for instance the papers [9, 10] of Bratteli and Jorgensen. If we start with a contractive tuple $(T_1, \dots, T_n)$ satisfying $\sum T_i T_i^* = I$ and consider the minimal isometric dilation we actually have a representation of the Cuntz algebra. Very interesting results on classification of these representation up to unitary equivalence in terms of invariants determined by $(T_1, \dots, T_n)$ have been obtained by Davidson, Kribs, and Shpigel [14], where the operators T_i act on a finite dimensional space. It is a natural question as to what are the representations of Cuntz algebra one can get by dilating contractive tuples which are also commuting. Surprisingly they are very few and are all determined by the GNS representations of

the so-called Cuntz states. This result we obtain in Section 4, as an application of the main result. Unlike the work of Davidson and Kribs, we need not restrict the operators T_i to be acting on finite dimensional spaces.

All the Hilbert spaces we consider will be complex and separable. For a subspace $\mathcal{H}$ of a Hilbert space $P_{\mathcal{H}}$ will denote the orthogonal projection onto $\mathcal{H}$. For fixed $n \geq 2$, we need two standard n -tuples of operators, denoted by $\underline{V}$ and $\underline{S}$ acting on Fock spaces. For any Hilbert space $\mathcal{K}$, we have the full Fock space over $\mathcal{K}$ denoted by $\Gamma(\mathcal{K})$ and the Boson (or symmetric) Fock space over $\mathcal{K}$ denoted by $\Gamma_s(\mathcal{K})$ as,

$$\Gamma(\mathcal{K}) = \mathbb{C} \oplus \mathcal{K} \oplus \mathcal{K}^{\otimes 2} \oplus \cdots \oplus \mathcal{K}^{\otimes m} \oplus \cdots,$$

$$\Gamma_s(\mathcal{K}) = \mathbb{C} \oplus \mathcal{K} \oplus \mathcal{K}^{\otimes^2} \oplus \cdots \oplus \mathcal{K}^{\otimes^m} \oplus \cdots,$$

where $\mathcal{K}^{\otimes^m}$ denotes m -fold symmetric tensor product. We will consider the Boson Fock space as a subspace of the full Fock space in the natural way. We denote the vacuum vector $1 \oplus 0 \oplus \cdots$ (in either of Fock spaces) by ω . Let $\mathbb{C}^n$ be the n -dimensional complex Euclidian space with usual inner product and $\Gamma(\mathbb{C}^n)$ be the full Fock space over $\mathbb{C}^n$. Let $\{e_1, \dots, e_n\}$ be the standard ortho-normal basis of $\mathbb{C}^n$. Then the (left) creation operators V_i on $\Gamma(\mathbb{C}^n)$ are defined by

$$V_i x = e_i \otimes x$$

where $1 \leq i \leq n$ and $x \in \Gamma(\mathbb{C}^n)$ (Of course, here $e_i \otimes \omega$ is interpreted as e_i). It is obvious that the tuple $\underline{V} = (V_1, \dots, V_n)$ consists of isometries with orthogonal ranges and it is contractive, in fact $\sum V_i V_i^* = I - E_0$, where E_0 is the projection on to the vacuum space. Let $\underline{S} = (S_1, \dots, S_n)$ be the tuple of operators on $\Gamma_s(\mathbb{C}^n)$ where, S_i is the compression of V_i to $\Gamma_s(\mathbb{C}^n)$:

$$S_i = P_{\Gamma_s(\mathbb{C}^n)} V_i|_{\Gamma_s(\mathbb{C}^n)}.$$

Clearly each V_i^* leaves $\Gamma_s(\mathbb{C}^n)$ invariant. Therefore, $S_i^* x = V_i^* x$, for $x \in \Gamma_s(\mathbb{C}^n)$. Then it is easy to see that $(S_1, \dots, S_n)$ is also a contractive tuple satisfying $\sum S_i S_i^* = I^s - E_0^s$ (where now I^s, E_0^s are identity, projection onto vacuum space in $\Gamma_s(\mathbb{C}^n)$). Moreover a simple computation shows that $S_i S_j = S_j S_i$ for all $1 \leq i, j \leq n$.

For operator tuples $(T_1, \dots, T_n)$ quite often we need to consider products of the form $T_{\alpha_1} T_{\alpha_2} \cdots T_{\alpha_m}$, where each $\alpha_k \in \{1, 2, \dots, n\}$. So it is convenient to have a notation for such products. Let Λ denote the set $\{1, 2, \dots, n\}$ and Λ^m denote the m -fold cartesian product of Λ for $m \geq 1$. Given $\alpha = (\alpha_1, \dots, \alpha_m)$ in Λ^m , $\underline{T}^\alpha$ will mean the operator $T_{\alpha_1} T_{\alpha_2} \cdots T_{\alpha_m}$. Let $\tilde{\Lambda}$ denote $\cup_{n=0}^\infty \Lambda^n$, where Λ^0 is just the set $\{0\}$ by convention and by $\underline{T}^0$ we would mean the identity operator of the Hilbert space where the operators T_i are acting. In a similar fashion for $\alpha \in \tilde{\Lambda}$, e^α will denote the vector $e_{\alpha_1} \otimes e_{\alpha_2} \otimes \cdots \otimes e_{\alpha_m}$ in the full Fock space $\Gamma(\mathbb{C}^n)$ and e^0 is the vacuum ω .

2. MAXIMAL COMMUTING PIECE AND DILATION

DEFINITION 2. Let $\mathcal{H}, \mathcal{L}$ be two Hilbert spaces such that $\mathcal{H}$ is a closed subspace of $\mathcal{L}$. Suppose $\underline{T}, \underline{R}$ are n -tuples of bounded operators on $\mathcal{H}, \mathcal{L}$ respectively. Then $\underline{R}$ is called a *dilation* of $\underline{T}$ if

$$R_i^* u = T_i^* u$$

for all $u \in \mathcal{H}$, $1 \leq i \leq n$. In such a case $\underline{T}$ is called a *piece* of $\underline{R}$. If further $\underline{T}$ is a commuting tuple (i. e., $T_i T_j = T_j T_i$, for all i, j), then it is called a commuting piece of $\underline{R}$. A dilation $\underline{R}$ of $\underline{T}$ is said to be a *minimal dilation* if $\overline{\text{span}}\{\underline{R}^\alpha h : \alpha \in \tilde{\Lambda}, h \in \mathcal{H}\} = \mathcal{L}$.

In this Definition we note that if $\underline{R}$ is a dilation of $\underline{T}$, then $\mathcal{H}$ is a co-invariant subspace of $\underline{R}$, that is, it is left invariant by all R_i^* . It is standard (see [18]) to call $(R_1^*, \dots, R_n^*)$ as an *extension* of $(T_1^*, \dots, T_n^*)$ and $(T_1^*, \dots, T_n^*)$ as a *part* of $(R_1^*, \dots, R_n^*)$. In such a situation it is easy to see that for any $\alpha, \beta \in \tilde{\Lambda}$, $\underline{T}^\alpha(\underline{T}^\beta)^*$ is the compression of $\underline{R}^\alpha(\underline{R}^\beta)^*$ to $\mathcal{H}$, that is,

$$(2.1) \quad \underline{T}^\alpha(\underline{T}^\beta)^* = P_{\mathcal{H}} \underline{R}^\alpha(\underline{R}^\beta)^*|_{\mathcal{H}}.$$

We may extend this relation to any polynomials p, q in n -noncommuting variables to have

$$p(\underline{T})(q(\underline{T}))^* = P_{\mathcal{H}} p(\underline{R})(q(\underline{R}))^*|_{\mathcal{H}}.$$

Usually it is property (2.1) is all that one demands of a dilation. But we have imposed a condition of co-invariance in Definition 2, as it is very convenient to have it this way for our purposes.

Now we look at commuting pieces of tuples. Let $\underline{R}$ be a n -tuple of bounded operators on a Hilbert space $\mathcal{L}$. Consider

$$\mathcal{C}(\underline{R}) = \{\mathcal{M} : \mathcal{M} \text{ is a co-invariant subspace for each } R_i, R_i^* R_j^* h = R_j^* R_i^* h, \forall h \in \mathcal{M}, \forall i, j\}.$$

So $\mathcal{C}(\underline{R})$ consists of all co-invariant subspaces of a n -tuple of operators $\underline{R}$ such that the compressions form a commuting tuple. It is a complete lattice, in the sense that arbitrary intersections and closures of arbitrary unions of such spaces are again in this collection. Therefore it has a maximal element. We denote it by $\mathcal{L}^c(\underline{R})$ (or by $\mathcal{L}^c$ when the tuple under consideration is clear).

DEFINITION 3. Suppose $\underline{R}$ is a n -tuple of operators on a Hilbert space $\mathcal{L}$. Then the *maximal commuting piece* of $\underline{R}$ is defined as the commuting piece $\underline{R}^c = (R_1^c, \dots, R_n^c)$ obtained by compressing $\underline{R}$ to the maximal element $\mathcal{L}^c(\underline{R})$ of $\mathcal{C}(\underline{R})$. The maximal commuting piece is said to be *trivial* if the space $\mathcal{L}^c(\underline{R})$ is just the zero space.

It is quite easy to get tuples with trivial commuting piece, as tuples with no non-trivial co-invariant subspaces have this property. Of course, our main interest lies in tuples with non-trivial commuting pieces. The following result is quite useful in determining the maximal commuting piece.

PROPOSITION 4. Let $\underline{R}$ be a n -tuple of bounded operators on a Hilbert space $\mathcal{L}$. Let $\mathcal{K}_{ij} = \overline{\text{span}}\{\underline{R}^\alpha(R_i R_j - R_j R_i)h : h \in \mathcal{L}, \alpha \in \tilde{\Lambda}\}$ for all $1 \leq i, j \leq n$, and $\mathcal{K} = \overline{\text{span}}\{\cup_{i,j=1}^n \mathcal{K}_{ij}\}$. Then $\mathcal{L}^c(\underline{R}) = \mathcal{K}^\perp$. In other words, $\mathcal{L}^c(\underline{R}) = \{h \in \mathcal{L} : (R_i^* R_j^* - R_j^* R_i^*)(\underline{R}^\alpha)^* h = 0, \forall 1 \leq i, j \leq n, \alpha \in \tilde{\Lambda}\}$.

PROOF: Firstly $\mathcal{K}^\perp$ is a co-invariant subspace of $\underline{R}$ is obvious as each R_i leaves $\mathcal{K}$ invariant. Now for $i, j \in \{1, 2, \dots, n\}$, and $h_1 \in \mathcal{L}^c, h_2 \in \mathcal{L}$,

$$\langle (R_i^* R_j^* - R_j^* R_i^*) h_1, h_2 \rangle = \langle h_1, (R_j R_i - R_i R_j) h_2 \rangle = 0.$$

So we get $(R_i^* R_j^* - R_j^* R_i^*) h_1 = 0$. Now if $\mathcal{M}$ is an element of $\mathcal{C}(\underline{R})$, take $i, j \in \{1, \dots, n\}, \alpha \in \tilde{\Lambda}, h_1 \in \mathcal{M}, h \in \mathcal{L}$. We have

$$\langle h_1, \underline{R}^\alpha(R_i R_j - R_j R_i) h \rangle = \langle (R_j^* R_i^* - R_i^* R_j^*)(\underline{R}^\alpha)^* h_1, h \rangle = 0$$

as $(\underline{R}^\alpha)^* h_1 \in \mathcal{M}$. Hence $\mathcal{M}$ is contained in $\mathcal{K}^\perp$. Now the last statement is easy to see. $\diamond$

COROLLARY 5. *Suppose $\underline{R}, \underline{T}$ are n -tuples of operators on two Hilbert spaces $\mathcal{L}, \mathcal{M}$. Then the maximal commuting piece of $(R_1 \oplus T_1, \dots, R_n \oplus T_n)$ acting on $\mathcal{L} \oplus \mathcal{M}$ is $(R_1^c \oplus T_1^c, \dots, R_n^c \oplus T_n^c)$ acting on $\mathcal{L}^c \oplus \mathcal{M}^c$. The maximal commuting piece of $(R_1 \otimes I, \dots, R_n \otimes I)$ acting on $\mathcal{L} \otimes \mathcal{M}$ is $(R_1^c \otimes I, \dots, R_n^c \otimes I)$ acting on $\mathcal{L}^c \otimes \mathcal{M}$.*

PROOF: Clear from Proposition 4. $\diamond$

PROPOSITION 6. *Let $\underline{V} = (V_1, \dots, V_n)$ and $\underline{S} = (S_1, \dots, S_n)$ be standard contractive tuples on full Fock space $\Gamma(\mathbb{C}^n)$ and Boson Fock space $\Gamma_s(\mathbb{C}^n)$ respectively. Then the maximal commuting piece of $\underline{V}$ is $\underline{S}$.*

PROOF: As we have already noted in the Introduction, $\underline{S}$ is a commuting piece of $\underline{V}$. To show maximality we make use of Proposition 4. Suppose $x \in \Gamma(\mathbb{C}^n)$ and $\langle x, \underline{V}^\alpha (V_j V_i - V_i V_j) y \rangle = 0$ for all $\alpha \in \tilde{\Lambda}, 1 \leq i, j \leq n$ and $y \in \Gamma(\mathbb{C}^n)$. We wish to show that $x \in \Gamma_s(\mathbb{C}^n)$. Suppose x_m is the m -particle component of x , that is, $x = \bigoplus_{m \geq 0} x_m$ with $x_m \in (\mathbb{C}^n)^{\otimes m}$ for $m \geq 0$. For $m \geq 2$ and any permutation σ of $\{1, 2, \dots, m\}$ we need to show that the unitary $U_\sigma : (\mathbb{C}^n)^{\otimes m} \rightarrow (\mathbb{C}^n)^{\otimes m}$, defined by

$$U_\sigma(u_1 \otimes \dots \otimes u_m) = u_{\sigma^{-1}(1)} \otimes \dots \otimes u_{\sigma^{-1}(m)},$$

leaves x_m fixed. Since the group of permutations of $\{1, 2, \dots, m\}$ is generated by permutations $\{(1, 2), \dots, (m-1, m)\}$ it is enough to verify $U_\sigma(x_m) = x_m$ for permutations σ of the form $(i, i+1)$. So fix m and i with $m \geq 2$ and $1 \leq i \leq (m-1)$. We have

$$\langle \bigoplus_p x_p, \underline{V}^\alpha (V_k V_l - V_l V_k) y \rangle = 0,$$

for every $y \in \Gamma(\mathbb{C}^n)$, $1 \leq k, l \leq n$. As α is arbitrary, this means that

$$\langle x_m, z \otimes (e_k \otimes e_l - e_l \otimes e_k) \otimes w \rangle = 0$$

for any $z \in (\mathbb{C}^n)^{\otimes(i-1)}, w \in (\mathbb{C}^n)^{\otimes(m-i-1)}$. This clearly implies $U_\sigma(x_m) = x_m$, for $\sigma = (i, i+1)$. $\diamond$

Now let us see how the maximal commuting piece behaves with respect to the operation of taking dilations. Before considering specific dilations we have the following general statement.

PROPOSITION 7. *Suppose $\underline{T}, \underline{R}$ are n -tuples of bounded operators on $\mathcal{H}, \mathcal{L}$, with $\mathcal{H} \subseteq \mathcal{L}$, such that $\underline{R}$ is a dilation of $\underline{T}$. Then $\mathcal{H}^c(\underline{T}) = \mathcal{L}^c(\underline{R}) \cap \mathcal{H}$ and $\underline{R}^c$ is a dilation of $\underline{T}^c$.*

PROOF: We have $R_i^* h = T_i^* h$, for $h \in \mathcal{H}$. Therefore, $(R_i^* R_j^* - R_j^* R_i^*)(\underline{R}^\alpha)^* h = (T_i^* T_j^* - T_j^* T_i^*)(\underline{T}^\alpha)^* h$ for $h \in \mathcal{H}$, $1 \leq i, j \leq n$, and $\alpha \in \tilde{\Lambda}$. Now the first part of the result is clear from Proposition 4. Further for $h \in \mathcal{L}^c(\underline{R})$, $R_i^* h = (R_i^c)^* h$ and so for $h \in \mathcal{H}^c(\underline{T}) = \mathcal{L}^c(\underline{R}) \cap \mathcal{H}$, $(R_i^c)^* h = R_i^* h = T_i^* h = (T_i^c)^* h$. This proves the claim. $\diamond$

DEFINITION 8. Let $\underline{T} = (T_1, \dots, T_n)$ be a contractive tuple on a Hilbert space $\mathcal{H}$. The operator $\Delta_{\underline{T}} = [I - (T_1 T_1^* + \dots + T_n T_n^*)]^{1/2}$ is called the *defect operator* of $\underline{T}$ and the subspace $\overline{\Delta_{\underline{T}}(\mathcal{H})}$ is called the *defect space* of $\underline{T}$. The tuple $\underline{T}$ is said to be *pure* if $\sum_{\alpha \in \Lambda^m} \underline{T}^\alpha (\underline{T}^\alpha)^*$ converges to zero in strong operator topology as m tends to infinity.

Suppose $\sum T_i T_i^* = I$, then it is easy to see that $\sum_{\alpha \in \Lambda^m} \underline{T}^\alpha (\underline{T}^\alpha)^* = I$ for all m and there is no-way this sequence can converge to zero. So in the pure case the defect operator and the defect spaces are non-trivial.

First we restrict our attention to pure tuples. The reason for this is that it is very easy to write down standard dilations for pure tuples. So let $\mathcal{H}$ be a complex, separable Hilbert space and let $\underline{T}$ be a pure contractive tuple on $\mathcal{H}$. Take $\tilde{\mathcal{H}} = \Gamma(\mathbb{C}^n) \otimes \overline{\Delta_{\underline{T}}(\mathcal{H})}$, and define an operator $A : \mathcal{H} \rightarrow \tilde{\mathcal{H}}$ by

$$(2.2) \quad Ah = \sum_{\alpha} e^{\alpha} \otimes \Delta_{\underline{T}}(\underline{T}^{\alpha})^* h,$$

where the sum is taken over all $\alpha \in \tilde{\Lambda}$. It is well-known ([23], [2]) and also easily verifiable using the pureness of $\underline{T}$, that A is an isometry with

$$A^*(e^{\alpha} \otimes h) = \underline{T}^{\alpha} \Delta_{\underline{T}} h \text{ for } \alpha \in \tilde{\Lambda}, h \in \overline{\Delta_{\underline{T}}(\mathcal{H})}.$$

Now $\mathcal{H}$ is considered as a subspace of $\tilde{\mathcal{H}}$ by identifying vectors $h \in \mathcal{H}$ with $Ah \in \tilde{\mathcal{H}}$. Then by noting that each $V_i^* \otimes I$ leaves the range of A invariant and $\underline{T}^{\alpha} = A^*(\underline{V}^{\alpha} \otimes I)A$ for all $\alpha \in \tilde{\Lambda}$ it is seen that the tuple $\tilde{\underline{V}} = (V_1 \otimes I, \dots, V_n \otimes I)$ of operators on $\tilde{\mathcal{H}}$ is a realization of the minimal isometric dilation of $\underline{T}$. Now if $\underline{T}$ is a commuting tuple, it is easy to see that the range of A is contained in $\tilde{\mathcal{H}}_s = \Gamma_s(\mathbb{C}^n) \otimes \overline{\Delta_{\underline{T}}(\mathcal{H})}$. In other words now $\mathcal{H}$ can be considered as a subspace of $\tilde{\mathcal{H}}_s$. Moreover, $\tilde{\underline{S}} = (S_1 \otimes I, \dots, S_n \otimes I)$, as a tuple of operators in $\tilde{\mathcal{H}}_s$ is a realization of the standard commuting dilation of $(T_1, \dots, T_n)$. More abstractly, if $\underline{T}$ is commuting and pure, the standard commuting dilation of it is got by embedding $\mathcal{H}$ isometrically in $\Gamma_s(\mathbb{C}^n) \otimes \mathcal{K}$, for some Hilbert space $\mathcal{K}$, such that $(S_1 \otimes I_{\mathcal{K}}, \dots, S_n \otimes I_{\mathcal{K}})$ is a dilation of $\underline{T}$ and $\overline{\text{span}}\{(\underline{S}^{\alpha} \otimes I_{\mathcal{K}})h : h \in \mathcal{H}, \alpha \in \tilde{\Lambda}\} = \Gamma_s(\mathbb{C}^n) \otimes \mathcal{K}$. Up to unitary equivalence there is unique such dilation and $\dim(\mathcal{K}) = \text{rank}(\Delta_{\underline{T}})$.

THEOREM 9. *Let $\underline{T}$ be a pure contractive tuple on a Hilbert space $\mathcal{H}$. Then the maximal commuting piece $\tilde{\underline{V}}^c$ of the minimal isometric dilation $\tilde{\underline{V}}$ of $\underline{T}$ is a realization of the standard commuting dilation of $\underline{T}^c$ if and only if $\Delta_{\underline{T}}(\mathcal{H}) = \overline{\Delta_{\underline{T}}(\mathcal{H}^c(\underline{T}))}$. In such a case $\text{rank}(\Delta_{\underline{T}}) = \text{rank}(\Delta_{\underline{T}^c}) = \text{rank}(\Delta_{\tilde{\underline{V}}}) = \text{rank}(\Delta_{\tilde{\underline{V}}^c})$.*

PROOF: We denote $\mathcal{H}^c(\underline{T})$, $\overline{\Delta_{\underline{T}}(\mathcal{H})}$, $\overline{\Delta_{\underline{T}}(\mathcal{H}^c(\underline{T}))}$ by $\mathcal{H}^c$, $\mathcal{M}$, and $\mathcal{M}^c$ respectively. It is obvious that $\underline{T}^c$ is also a pure contractive tuple. We already know from Proposition 7 that $\tilde{\underline{V}}^c = (\underline{S} \otimes I_{\mathcal{M}})$ on $\Gamma_s(\mathbb{C}^n) \otimes \mathcal{M}$ is a dilation of $\underline{T}^c$. It is the standard dilation if and only if $\mathcal{L} := \overline{\text{span}}\{(\underline{S}^{\alpha} \otimes I_{\mathcal{M}})Ah : h \in \mathcal{H}^c, \alpha \in \tilde{\Lambda}\}$ is equal to $\Gamma_s(\mathbb{C}^n) \otimes \mathcal{M}$, where $A : \mathcal{H} \rightarrow \tilde{\mathcal{H}}$ is the isometry defined by (2.2).

From the definition of A , using the commutativity of the operators T_i , it is clear that for $h \in \mathcal{H}^c$, $Ah \in \Gamma_s(\mathbb{C}^n) \otimes \mathcal{M}^c$. Hence $\mathcal{L} \subseteq \Gamma_s(\mathbb{C}^n) \otimes \mathcal{M}^c$. Further, as $(\underline{S} \otimes I_{\mathcal{M}})$ is a dilation, $(S_i^* \otimes I_{\mathcal{M}})$ leaves $A(\mathcal{H}^c)$ invariant. Therefore, $((I - \sum S_i S_i^*) \otimes I_{\mathcal{M}})Ah \in \mathcal{L}$ for $h \in \mathcal{H}^c$. But $(I - \sum S_i S_i^*)$ being the projection onto the vacuum space, $((I - \sum S_i S_i^*) \otimes I_{\mathcal{M}})Ah = \omega \otimes \Delta_{\underline{T}} h$. As $\{\underline{S}^{\alpha}, \alpha \in \tilde{\Lambda}\}$ spans whole of $\Gamma_s(\mathbb{C}^n)$ we get that $\Gamma_s(\mathbb{C}^n) \otimes \mathcal{M}^c \subseteq \mathcal{L}$. Hence $\mathcal{L} = \Gamma_s(\mathbb{C}^n) \otimes \mathcal{M}^c$ and this way we have proved the first claim.

Now suppose $\tilde{\underline{V}}^c$ is a realization of the standard commuting dilation of $\underline{T}^c$. This in particular means that $\text{rank}(\Delta_{\underline{T}^c}) = \text{rank}(\Delta_{\tilde{\underline{V}}^c})$. Also as $\tilde{\underline{V}}$ is the minimal isometric dilation of $\underline{T}$, $\text{rank}(\Delta_{\underline{T}}) = \text{rank}(\Delta_{\tilde{\underline{V}}})$. Further as $\tilde{\underline{V}}^c = (\underline{S} \otimes I_{\mathcal{M}})$, $\text{rank}(\Delta_{\tilde{\underline{V}}^c}) = \dim(\mathcal{M}) = \text{rank}(\Delta_{\underline{T}})$. $\diamond$

We may ask whether the equality of ranks in this Theorem is good enough to make a converse statement. To answer this we make use of the following simple lemma.

LEMMA 10. *Suppose*

$$M = \begin{bmatrix} A & B^* \\ B & C \end{bmatrix}$$

is a bounded positive operator on some Hilbert space. Then $\text{rank}(A) = \text{rank} \left(\begin{bmatrix} A \\ B \end{bmatrix} \right)$

PROOF: Without loss of generality we can assume that M is a contraction. Then it is a folklore theorem that there exists a contraction D such that $B = C^{\frac{1}{2}}DA^{\frac{1}{2}}$. Now

$$\begin{bmatrix} A \\ B \end{bmatrix} = \begin{bmatrix} A^{\frac{1}{2}} \\ C^{\frac{1}{2}}D \end{bmatrix} \begin{bmatrix} A^{\frac{1}{2}} \end{bmatrix},$$

and hence $\text{rank} \left(\begin{bmatrix} A \\ B \end{bmatrix} \right) \leq \text{rank}(A^{\frac{1}{2}})$. But A being positive, $\text{rank}(A) = \text{rank}(A^{\frac{1}{2}})$. Therefore $\text{rank} \left(\begin{bmatrix} A \\ B \end{bmatrix} \right) \leq \text{rank}(A) \leq \text{rank} \left(\begin{bmatrix} A \\ B \end{bmatrix} \right)$. $\diamond$

REMARK 11. *Let $\underline{T}$ be a pure contractive tuple on a Hilbert space $\mathcal{H}$ with minimal isometric dilation $\tilde{\underline{V}}$. If $\text{rank } \Delta_{\underline{T}}$ and $\text{rank } \Delta_{\underline{T}^c}$ are finite and equal then $\tilde{\underline{V}}^c$ is a realization of the standard commuting dilation of $\underline{T}^c$.*

PROOF: In view of Theorem 9 we need to show that $\overline{\Delta_{\underline{T}}(\mathcal{H})} = \overline{\Delta_{\underline{T}}(\mathcal{H}^c(\underline{T}))}$. Since $\overline{\Delta_{\underline{T}}(\mathcal{H})} \supseteq \overline{\Delta_{\underline{T}}(\mathcal{H}^c(\underline{T}))}$, and these spaces are now finite dimensional, it suffices to show that their dimensions are equal or $\text{rank}(\Delta_{\underline{T}}) = \text{rank}(\Delta_{\underline{T}}P_{\mathcal{H}^c})$. Clearly $\text{rank}(\Delta_{\underline{T}}) \geq \text{rank}(\Delta_{\underline{T}}P_{\mathcal{H}^c})$. Also by assumption, $\text{rank}(\Delta_{\underline{T}}) = \text{rank}(\Delta_{\underline{T}^c})$. By positivity $\text{rank}(\Delta_{\underline{T}^c}) = \text{rank}(\Delta_{\underline{T}^c}^2)$. And then by previous Lemma $\text{rank}(\Delta_{\underline{T}^c}^2) = \text{rank}(P_{\mathcal{H}^c}(\Delta_{\underline{T}}^2)P_{\mathcal{H}^c}) = \text{rank}(\Delta_{\underline{T}}^2P_{\mathcal{H}^c}) \leq \text{rank}(\Delta_{\underline{T}}P_{\mathcal{H}^c})$. $\diamond$

If both the ranks are infinite then we can not ensure that $\overline{\Delta_{\underline{T}}(\mathcal{H})} = \overline{\Delta_{\underline{T}}(\mathcal{H}^c(\underline{T}))}$ is seen by the following example.

EXAMPLE 12. Let $\underline{R} = (R_1, R_2)$ be a commuting pure contractive 2-tuple on an infinite dimensional Hilbert space $\mathcal{H}_0$ (We can even take R_1, R_2 as scalars) such that $\overline{\Delta_{\underline{R}}(\mathcal{H}_0)}$ is infinite dimensional. Take $\mathcal{H} = \mathcal{H}_0 \oplus \mathbb{C}^2$, and let T_1, T_2 be operators on $\mathcal{H}$ defined by

$$T_1 = \begin{bmatrix} R_1 & & \\ & 0 & t_1 \\ & 0 & 0 \end{bmatrix}, T_2 = \begin{bmatrix} R_2 & & \\ & 0 & 0 \\ & t_2 & 0 \end{bmatrix},$$

where t_1, t_2 are any two scalars, $0 < t_1, t_2 < 1$. Then $\underline{T} = (T_1, T_2)$ is a pure contractive tuple. Making use of Corollary 5, $\mathcal{H}^c(\underline{T}) = \mathcal{H}_0$ (thought of as a subspace of $\mathcal{H}$ in the natural way) and the maximal commuting piece of $\underline{T}$ is (R_1, R_2) , and therefore $\text{rank}(\Delta_{\underline{T}^c}) = \text{rank}(\Delta_{\underline{T}}) = \infty$. But $\overline{\Delta_{\underline{T}}(\mathcal{H})} = \overline{\Delta_{\underline{R}}(\mathcal{H}_0)} \oplus \mathbb{C}^2$.

We do not know how to extend Theorem 9 to contractive tuples which are not necessarily pure.

3. COMMUTING TUPLES

In this Section we wish to consider commutative contractive tuples. Let us begin with describing the way one obtains two standard dilations for such tuples.

Recall standard tuples $\underline{V}$ and $\underline{S}$ on Fock spaces $\Gamma(\mathbb{C}^n)$, and $\Gamma_s(\mathbb{C}^n)$ respectively, introduced in the Introduction. Let $C^*(\underline{V})$, and $C^*(\underline{S})$ be unital C^* algebras generated by them. For any $\alpha, \beta \in \tilde{\Lambda}$, $\underline{V}^\alpha(I - \sum V_i V_i^*)(\underline{V}^\beta)^*$ is the rank one operator $x \mapsto \langle e^\beta, x \rangle e^\alpha$, formed by basis vectors e^α, e^β . So $C^*(\underline{V})$ contains all compact operators. In a similar way we see that $C^*(\underline{S})$ also contains all compact operators of $\Gamma_s(\mathbb{C}^n)$. As $V_i^* V_j = \delta_{ij}$, it is easy to see that $C^*(\underline{V}) = \overline{\text{span}} \{ \underline{V}^\alpha (\underline{V}^\beta)^* : \alpha, \beta \in \tilde{\Lambda} \}$. By explicit computation commutators $[S_i^*, S_j]$ are compact for all i, j (See [5], Proposition 5.3, or [8]). Therefore we can also obtain $C^*(\underline{S}) = \overline{\text{span}} \{ \underline{S}^\alpha (\underline{S}^\beta)^* : \alpha, \beta \in \tilde{\Lambda} \}$.

Suppose $\underline{T}$ is a contractive tuple on a Hilbert space $\mathcal{H}$. We obtain a certain completely positive map (Popescu's Poisson transform) from $C^*(\underline{V})$ to $\mathcal{B}(\mathcal{H})$, as follows. For $0 < r < 1$ the tuple $r\underline{T} = (rT_1, \dots, rT_n)$ is clearly a pure contraction. So by (2.2) we have an isometry $A_r : \mathcal{H} \rightarrow \Gamma(\mathbb{C}^n) \otimes \Delta_r(\mathcal{H})$ defined by

$$A_r h = \sum_{\alpha} e^{\alpha} \otimes \Delta_r((r\underline{T})^{\alpha})^* h, \quad h \in \mathcal{H},$$

where $\Delta_r = (I - r^2 \sum T_i T_i^*)^{\frac{1}{2}}$. So for every $0 < r < 1$ we have a completely positive map $\psi_r : C^*(\underline{V}) \rightarrow \mathcal{B}(\mathcal{H})$ defined by

$$\psi_r(X) = A_r^*(X \otimes I)A_r, \quad X \in C^*(\underline{V}).$$

By taking limit as r increases to 1 (See [23] or [2] for details), we obtain a unital completely positive map ψ from $C^*(\underline{V})$ to $\mathcal{B}(\mathcal{H})$ satisfying

$$\psi(\underline{V}^{\alpha}(\underline{V}^{\beta})^*) = \underline{T}^{\alpha}(\underline{T}^{\beta})^* \text{ for } \alpha, \beta \in \tilde{\Lambda}.$$

As $C^*(\underline{V}) = \overline{\text{span}} \{ \underline{V}^{\alpha}(\underline{V}^{\beta})^* : \alpha, \beta \in \tilde{\Lambda} \}$, ψ is the unique such completely positive map. Now consider the minimal Stinespring dilation of ψ . So we have a Hilbert space $\tilde{\mathcal{H}}$ containing $\mathcal{H}$, and a unital $*$ -homomorphism $\pi : C^*(\underline{V}) \rightarrow \mathcal{B}(\tilde{\mathcal{H}})$, such that

$$\psi(X) = P_{\mathcal{H}}\pi(X)|_{\mathcal{H}} \quad \forall X \in C^*(\underline{V}),$$

and $\overline{\text{span}} \{ \pi(X)h : X \in C^*(\underline{V}), h \in \mathcal{H} \} = \tilde{\mathcal{H}}$. Taking $\tilde{\underline{V}} = (\tilde{V}_1, \dots, \tilde{V}_n) = (\pi(V_1), \dots, \pi(V_n))$, one verifies that each $(\tilde{V}_i)^*$ leaves $\mathcal{H}$ invariant and $\tilde{\underline{V}}$ is the unique minimal isometric dilation of $\underline{V}$.

In a similar fashion if $\underline{T}$ is commuting by considering $C^*(\underline{S})$ instead of $C^*(\underline{V})$, and restricting A_r in the range to $\Gamma_s(\mathbb{C}^n)$, and taking limits as before (See [5], [23], [2]) we obtain the unique unital completely positive map $\phi : C^*(\underline{S}) \rightarrow \mathcal{B}(\mathcal{H})$, satisfying

$$\phi(\underline{S}^{\alpha}(\underline{S}^{\beta})^*) = \underline{T}^{\alpha}(\underline{T}^{\beta})^* \quad \alpha, \beta \in \tilde{\Lambda}.$$

Consider the minimal Stinespring dilation of ϕ . Here we have a Hilbert space $\mathcal{H}_1$ containing $\mathcal{H}$ and a unital $*$ -homomorphism $\pi_1 : C^*(\underline{S}) \rightarrow \mathcal{B}(\mathcal{H}_1)$, such that

$$\phi(X) = P_{\mathcal{H}}\pi_1(X)|_{\mathcal{H}} \quad \forall X \in C^*(\underline{S}),$$

and $\overline{\text{span}} \{ \pi_1(X)h : X \in C^*(\underline{S}), h \in \mathcal{H} \} = \mathcal{H}_1$. Taking $\tilde{\underline{S}} = (\tilde{S}_1, \dots, \tilde{S}_n) = (\pi_1(S_1), \dots, \pi_1(S_n))$, $\tilde{\underline{S}}$ is the standard commuting dilation of $\underline{T}$ by definition (It is not difficult to verify that it is a

minimal dilation in the sense of our Definition 2). As minimal Stinespring dilation is unique up to unitary equivalence, standard commuting dilation is also unique up to unitary equivalence.

THEOREM 13. (*Main Theorem*) Suppose $\underline{T}$ is a commuting contractive tuple on a Hilbert space $\mathcal{H}$. Then the maximal commuting piece of the minimal isometric dilation of $\underline{T}$ is a realization of the standard commuting dilation of $\underline{T}$.

Our approach to prove this theorem is as follows. First we consider the standard commuting dilation of $\underline{T}$ on a Hilbert space $\mathcal{H}_1$ as described above. Now the standard tuple $\underline{S}$ is also a contractive tuple. So we have a unique unital completely positive map $\eta : C^*(\underline{V}) \rightarrow C^*(\underline{S})$, satisfying

$$\eta(\underline{V}^\alpha(\underline{V}^\beta)^*) = \underline{S}^\alpha(\underline{S}^\beta)^* \quad \alpha, \beta \in \tilde{\Lambda}.$$

Now clearly $\psi = \phi \circ \eta$. Consider the minimal Stinespring dilation of the composed map $\pi_1 \circ \eta : C^*(\underline{V}) \rightarrow \mathcal{B}(\mathcal{H}_1)$. Here we obtain a Hilbert space $\mathcal{H}_2$ containing $\mathcal{H}_1$ and a unital $*$ -homomorphism $\pi_2 : C^*(\underline{V}) \rightarrow \mathcal{B}(\mathcal{H}_2)$, such that

$$\pi_1 \circ \eta(X) = P_{\mathcal{H}_1} \pi_2(X)|_{\mathcal{H}_1}, \quad \forall X \in C^*(\underline{V}),$$

and $\overline{\text{span}} \{ \pi_2(X)h : X \in C^*(\underline{V}), h \in \mathcal{H}_1 \} = \mathcal{H}_2$. Now we have a commuting diagram as follows

$$\begin{array}{ccccc} & & & & \mathcal{B}(\mathcal{H}_2) \\ & & & & \downarrow \\ & & & & \mathcal{B}(\mathcal{H}_1) \\ & & & & \downarrow \\ & & & & \mathcal{B}(\mathcal{H}) \\ C^*(\underline{V}) & \xrightarrow{\eta} & C^*(\underline{S}) & \xrightarrow{\phi} & \mathcal{B}(\mathcal{H}) \\ & \nearrow \pi_2 & \nearrow \pi_1 & & \\ & & & & \end{array}$$

where all the down arrows are compression maps, horizontal arrows are unital completely positive maps and diagonal arrows are unital $*$ -homomorphisms.

Taking $\hat{\underline{V}} = (\hat{V}_1, \dots, \hat{V}_n) = (\pi_2(V_1), \dots, \pi_2(V_n))$, we need to show (i) $\hat{\underline{V}}$ is the minimal isometric dilation of $\underline{T}$ and (ii) $\hat{\underline{S}} = (\pi_1(S_1), \dots, \pi_1(S_n))$ is the maximal commuting piece of $\hat{\underline{V}}$. Due to uniqueness up to unitary equivalence of minimal Stinespring dilation, we have (i) if we can show that π_2 is a minimal dilation of $\psi = \phi \circ \eta$. For proving this we actually make use of (ii). At first we prove (ii) in a very special case.

DEFINITION 14. A n -tuple $\underline{T} = (T_1, \dots, T_n)$ of operators on a Hilbert space $\mathcal{H}$ is called a spherical unitary if it is commuting, each T_i is normal, and $T_1 T_1^* + \dots + T_n T_n^* = I$.

Actually, if $\mathcal{H}$ is a finite dimensional Hilbert space and $\underline{T}$ is a commuting tuple on $\mathcal{H}$ satisfying $\sum T_i T_i^* = I$, then it is automatically a spherical unitary, that is, each T_i is normal. This is the case because here standard commuting dilation of $\underline{T}$ is a tuple of normal operators and hence each T_i^* is subnormal (or see [6] for this result) and all finite dimensional subnormal operators are normal (see [18]).

Note that if $\underline{T}$ is a spherical unitary we have $\phi(\underline{S}^\alpha(I - \sum S_i S_i^*)(\underline{S}^\beta)^*) = \underline{T}^\alpha(I - \sum T_i T_i^*)(\underline{T}^\beta)^* = 0$ for any $\alpha, \beta \in \tilde{\Lambda}$. This forces that $\phi(X) = 0$ for any compact operator X in $C^*(\underline{S})$. Now as the

commutators $[S_i^*, S_j]$ are all compact we see that ϕ is a unital $*$ -homomorphism. So the minimal Stinespring dilation of ϕ is itself. So the following result yields Theorem 13 for spherical unitaries.

THEOREM 15. *Let $\underline{T}$ be a spherical unitary on a Hilbert space $\mathcal{H}$. Then the maximal commuting piece of the minimal isometric dilation of $\underline{T}$ is $\underline{T}$.*

As proof of this Theorem involves some lengthy computations we prefer to postpone it. But assuming this, we prove the main Theorem.

PROOF OF THEOREM 13 : As $C^*(\underline{S})$ contains the ideal of all compact operators by standard C^* -algebra theory we have a direct sum decomposition of π_1 as follows. Take $\mathcal{H}_1 = \mathcal{H}_{1C} \oplus \mathcal{H}_{1N}$ where $\mathcal{H}_{1C} = \overline{\text{span}}\{\pi(X)h : h \in \mathcal{H}, X \in C^*(\underline{S}) \text{ and } X \text{ is compact}\}$ and $\mathcal{H}_{1N} = \mathcal{H}_1 \ominus \mathcal{H}_{1C}$, Clearly $\mathcal{H}_{1C}$ is a reducing subspace for π_1 . Therefore

$$\pi_1(X) = \begin{pmatrix} \pi_{1C}(X) & \\ & \pi_{1N}(X) \end{pmatrix}$$

that is, $\pi_1 = \pi_{1C} \oplus \pi_{1N}$ where $\pi_{1C}(X) = P_{\mathcal{H}_{1C}} \pi_1(X) P_{\mathcal{H}_{1C}}$, $\pi_{1N}(X) = P_{\mathcal{H}_{1N}} \pi_1(X) P_{\mathcal{H}_{1N}}$. As observed by Arveson [5], $\pi_{1C}(X)$ is just the identity representation with some multiplicity. More precisely, $\mathcal{H}_{1C}$ can be factored as $\mathcal{H}_{1C} = \Gamma_s(\mathbb{C}^n) \otimes \overline{\Delta_{\underline{T}}(\mathcal{H})}$, such that $\pi_{1C}(X) = X \otimes I$, in particular $\pi_{1C}(S_i) = S_i \otimes I$. Also $\pi_{1N}(X) = 0$ for compact X . Therefore, taking $Z_i = \pi_{1N}(S_i)$, $\underline{Z} = (Z_1, \dots, Z_n)$ is a spherical unitary.

Now as $\pi_1 \circ \eta = (\pi_{1C} \circ \eta) \oplus (\pi_{1N} \circ \eta)$ and the minimal Stinespring dilation of a direct sum of two completely positive maps is the direct sum of minimal Stinespring dilations. So $\mathcal{H}_2$ decomposes as $\mathcal{H}_2 = \mathcal{H}_{2C} \oplus \mathcal{H}_{2N}$, where $\mathcal{H}_{2C}, \mathcal{H}_{2N}$ are orthogonal reducing subspaces of π_2 , such that π_2 also decomposes, say $\pi_2 = \pi_{2C} \oplus \pi_{2N}$, with

$$\pi_{1C} \circ \eta(X) = P_{\mathcal{H}_{1C}} \pi_{2C}(X)|_{\mathcal{H}_{1C}}, \quad \pi_{1N} \circ \eta(X) = P_{\mathcal{H}_{1N}} \pi_{2N}(X)|_{\mathcal{H}_{1N}},$$

for $X \in C^*(\underline{V})$ with $\mathcal{H}_{2C} = \overline{\text{span}}\{\pi_{2C}(X)h : X \in C^*(\underline{V}), h \in \mathcal{H}_{1C}\}$ and $\mathcal{H}_{2N} = \overline{\text{span}}\{\pi_{2N}(X)h : X \in C^*(\underline{V}), h \in \mathcal{H}_{1N}\}$. It is also not difficult to see that $\mathcal{H}_{2C} = \overline{\text{span}}\{\pi_{2C}(X)h : X \in C^*(\underline{V}), X \text{ compact}, h \in \mathcal{H}_{1C}\}$ and hence $\mathcal{H}_{2C}$ factors as $\mathcal{H}_{2C} = \Gamma(\mathbb{C}^n) \otimes \overline{\Delta_{\underline{T}}(\mathcal{H})}$ with $\pi_{2C}(V_i) = V_i \otimes I$. Also $(\pi_{2N}(V_1), \dots, \pi_{2N}(V_n))$ is a minimal isometric dilation of spherical isometry $(Z_1, \dots, Z_n)$. Now by Proposition 6, Theorem 15 and Corollary 5, we get that $(\pi_1(S_1), \dots, \pi_1(S_n))$ acting on $\mathcal{H}_1$ is the maximal commuting piece of $(\pi_2(V_1), \dots, \pi_2(V_n))$.

All that remains to show is that π_2 is the minimal Stinespring dilation of $\phi \circ \eta$. Suppose this is not the case. Then we get a reducing subspace $\mathcal{H}_{20}$ for π_2 by taking $\mathcal{H}_{20} = \overline{\text{span}}\{\pi_2(X)h : X \in C^*(\underline{V}), h \in \mathcal{H}\}$. Take $\mathcal{H}_{21} = \mathcal{H}_2 \ominus \mathcal{H}_{20}$ and correspondingly decompose π_2 as $\pi_2 = \pi_{20} \oplus \pi_{21}$,

$$\pi_2(X) = \begin{pmatrix} \pi_{20}(X) & \\ & \pi_{21}(X) \end{pmatrix}$$

Note that we already have $\mathcal{H} \subseteq \mathcal{H}_{20}$. We claim that $\mathcal{H}_2 \subseteq \mathcal{H}_{20}$. Firstly, as $\mathcal{H}_1$ is the space where the maximal commuting piece of $(\pi_2(V_1), \dots, \pi_2(V_n)) = (\pi_{20}(V_1) \oplus \pi_{21}(V_1), \dots, \pi_{20}(V_n) \oplus \pi_{21}(V_n))$ acts, by the first part of Corollary 5, $\mathcal{H}_1$ decomposes as $\mathcal{H}_1 = \mathcal{H}_{10} \oplus \mathcal{H}_{11}$ for some subspaces

$\mathcal{H}_{10} \subseteq \mathcal{H}_{20}$, and $\mathcal{H}_{11} \subseteq \mathcal{H}_{21}$. So for $X \in C^*(\underline{V})$, $P_{\mathcal{H}_1}\pi_2(X)P_{\mathcal{H}_1}$, has the form (see the diagram)

$$P_{\mathcal{H}_1}\pi_2(X)P_{\mathcal{H}_1} = \begin{pmatrix} \pi_{10} \circ \eta(X) & 0 & & \\ & 0 & 0 & \\ & & \pi_{11} \circ \eta(X) & 0 \\ & & 0 & 0 \end{pmatrix}$$

where π_{10}, π_{11} are compressions of π_1 to $\mathcal{H}_{10}, \mathcal{H}_{11}$ respectively. As the mapping η from $C^*(\underline{V})$ to $C^*(\underline{S})$ is clearly surjective, it follows that $\mathcal{H}_{10}, \mathcal{H}_{11}$ are reducing subspaces for π_1 . Now as $\mathcal{H}$ is contained in $\mathcal{H}_{20}$, in view of minimality of π_1 as a Stinespring dilation, $\mathcal{H}_1 \subseteq \mathcal{H}_{20}$. But then the minimality of π_2 shows that $\mathcal{H}_2 \subseteq \mathcal{H}_{20}$. Therefore, $\mathcal{H}_2 = \mathcal{H}_{20}$. $\diamond$

PROOF OF THEOREM 15 : Here we need a different presentation of the minimal isometric dilation. This is known as Schäffer construction [26] in the one variable case, and [20] is a good reference for the multivariate case. Here we decompose the dilation space $\tilde{\mathcal{H}}$ as $\tilde{\mathcal{H}} = \mathcal{H} \oplus (\Gamma(\mathbb{C}^n) \otimes \mathcal{D})$ where $\mathcal{D}$ is the closure of the range of operator

$$D : \underbrace{\mathcal{H} \oplus \cdots \oplus \mathcal{H}}_{n \text{ copies}} \rightarrow \underbrace{\mathcal{H} \oplus \cdots \oplus \mathcal{H}}_{n \text{ copies}}$$

and D is the positive square root of

$$D^2 = [\delta_{ij}I - T_i^*T_j]_{n \times n}.$$

Whenever it is convenient for us we identify $\underbrace{\mathcal{H} \oplus \cdots \oplus \mathcal{H}}_{n \text{ copies}}$ with $\mathbb{C}^n \otimes \mathcal{H}$ so that

$$(h_1, \dots, h_n) = \sum_{i=1}^n e_i \otimes h_i.$$

Then

$$(3.1) \quad D(h_1, \dots, h_n) = D\left(\sum_{i=1}^n e_i \otimes h_i\right) = \sum_{i=1}^n e_i \otimes \left(h_i - \sum_{j=1}^n T_i^*T_j h_j\right).$$

And the minimal isometric dilation $\tilde{V}_i$ has the form

$$(3.2) \quad \tilde{V}_i(h \oplus \sum_{\alpha \in \tilde{\Lambda}} e^\alpha \otimes d_\alpha) = T_i h \oplus D(e_i \otimes h) \oplus e_i \otimes \left(\sum_{\alpha \in \tilde{\Lambda}} e^\alpha \otimes d_\alpha\right)$$

for $h \in \mathcal{H}$, $d_\alpha \in \mathcal{D}$ for $\alpha \in \tilde{\Lambda}$, and $1 \leq i \leq n$ ($\mathbb{C}^n \omega \otimes \mathcal{D}$ has been identified with $\mathcal{D}$).

In the present case as $\sum T_i T_i^* = I$, by direct computation D^2 is seen to be a projection. So, D which is the positive square root of D^2 , is equal to D^2 . Also by Fuglede-Putnam theorem ([18], [25]), $\{T_1, \dots, T_n, T_1^*, \dots, T_n^*\}$ forms a commuting family of operators. Then we get

$$(3.3) \quad D(h_1, \dots, h_n) = \sum_{i,j=1}^n e_i \otimes T_j(T_j^* h_i - T_i^* h_j) = \sum_{i,j=1}^n e_i \otimes T_j(h_{ij})$$

where $h_{ij} = T_j^* h_i - T_i^* h_j$ for $1 \leq i, j \leq n$. Note that $h_{ii} = 0$ and $h_{ji} = -h_{ij}$.

Now we apply Proposition 4 to the tuple $\tilde{\underline{V}}$ acting on $\tilde{\mathcal{H}}$. Suppose $y \in \mathcal{H}^\perp \cap \tilde{\mathcal{H}}^c(\tilde{\underline{V}})$. We wish to show that $y = 0$. We assume $y \neq 0$ and arrive at a contradiction. One can decompose y as

$y = 0 \oplus \sum_{\alpha \in \tilde{\Lambda}} e^\alpha \otimes y_\alpha$, with $y_\alpha \in \mathcal{D}$. If for some α , $y_\alpha \neq 0$, then $\langle \omega \otimes y_\alpha, (\tilde{V}^\alpha)^* y \rangle = \langle e^\alpha \otimes y_\alpha, y \rangle = \langle y_\alpha, y_\alpha \rangle \neq 0$. Since each $(\tilde{V}_i)^*$ leaves $\tilde{\mathcal{H}}^c(\tilde{V})$ invariant, $(\tilde{V}^\alpha)^* y \in \tilde{\mathcal{H}}^c(\tilde{V})$. So without loss of generality we can assume $\|y_0\| = 1$.

Taking $\tilde{y}_m = \sum_{\alpha \in \Lambda^m} e^\alpha \otimes y_\alpha$, we get $y = 0 \oplus \oplus_{m \geq 0} (\tilde{y}_m)$. As $y_0 \in \mathcal{D}$, $y_0 = D(h_1, \dots, h_n)$, for some $(h_1, \dots, h_n)$ (Presently D being a projection its range is closed). Set $\tilde{x}_0 = \tilde{y}_0 = y_0$, and for $m \geq 1$,

$$\tilde{x}_m = \sum_{i_1, \dots, i_{m-1}, i, j=1}^n e_{i_1} \otimes \dots \otimes e_{i_{m-1}} \otimes e_i \otimes D(e_j \otimes T_{i_1}^* \dots T_{i_{m-1}}^* h_{ij}).$$

Clearly $\tilde{x}_m \in (\mathbb{C}^n)^{\otimes m} \otimes \mathcal{D}$ for all $m \in \mathbb{N}$. From the definition (3.2) of $\tilde{V}_i$, commutativity of the operators T_i , and the fact that D is projection, we have

$$\begin{aligned} \sum_{1 \leq i < j \leq n} (\tilde{V}_i \tilde{V}_j - \tilde{V}_j \tilde{V}_i) h_{ij} &= \sum_{1 \leq i < j \leq n} (T_i T_j h_{ij} - T_j T_i h_{ij}) \\ &+ \sum_{1 \leq i < j \leq n} D(e_i \otimes T_j h_{ij} - e_j \otimes T_i h_{ij}) \\ &+ \sum_{1 \leq i < j \leq n} (e_i \otimes D(e_j \otimes h_{ij}) - e_j \otimes D(e_i \otimes h_{ij})) \\ &= D \left\{ \sum_{1 \leq i < j \leq n} (e_i \otimes T_j h_{ij} - e_j \otimes T_i h_{ij}) \right\} + \sum_{i,j=1}^n e_i \otimes D(e_j \otimes h_{ij}) \\ &= D \left(\sum_{1 \leq i, j \leq n} e_i \otimes T_j h_{ij} \right) + \sum_{i,j=1}^n e_i \otimes D(e_j \otimes h_{ij}) \\ &= D^2(h_1, \dots, h_n) + \sum_{i,j=1}^n e_i \otimes D(e_j \otimes h_{ij}) \\ &= \tilde{x}_0 + \tilde{x}_1. \end{aligned}$$

Therefore $\langle y, \tilde{x}_0 + \tilde{x}_1 \rangle = 0$ by Proposition 4. Now for $m \geq 2$.

$$\begin{aligned} &\sum_{i_1, \dots, i_{m-1}=1}^n \tilde{V}_{i_1} \dots \tilde{V}_{i_{m-1}} \left(\sum_{i,j=1}^n (\tilde{V}_i \tilde{V}_j - \tilde{V}_j \tilde{V}_i) T_{i_1}^* \dots T_{i_{m-2}}^* T_j^* h_{i_{m-1}i} \right) \\ &= \sum_{i_1, \dots, i_{m-1}=1}^n \tilde{V}_{i_1} \dots \tilde{V}_{i_{m-1}} \left[\sum_{i,j=1}^n D(e_i \otimes T_j T_{i_1}^* \dots T_{i_{m-2}}^* T_j^* h_{i_{m-1}i} - e_j \otimes T_i T_{i_1}^* \dots T_{i_{m-2}}^* T_j^* h_{i_{m-1}i}) \right. \\ &\quad \left. + \sum_{i,j=1}^n \{ e_i \otimes D(e_j \otimes T_{i_1}^* \dots T_{i_{m-2}}^* T_j^* h_{i_{m-1}i}) - e_j \otimes D(e_i \otimes T_{i_1}^* \dots T_{i_{m-2}}^* T_j^* h_{i_{m-1}i}) \} \right] \end{aligned}$$

$$\begin{aligned}
&= \sum_{i_1, \dots, i_{m-1}=1}^n e_{i_1} \otimes \cdots \otimes e_{i_{m-1}} \otimes \\
&\quad [D(\sum_{i,j=1}^n e_i \otimes T_j T_{i_1}^* \cdots T_{i_{m-2}}^* T_j^* h_{i_{m-1}i} - e_j \otimes T_i T_{i_1}^* \cdots T_{i_{m-2}}^* T_j^* h_{i_{m-1}i}) \\
&\quad \{ \sum_{i,j=1}^n e_i \otimes D(e_j \otimes T_{i_1}^* \cdots T_{i_{m-2}}^* T_j^* h_{i_{m-1}i}) - \sum_{i,j=1}^n e_i \otimes D(e_j \otimes T_{i_1}^* \cdots T_{i_{m-2}}^* T_i^* h_{i_{m-1}j}) \}] \\
&\quad (\text{in the term above, } i \text{ and } j \text{ have been interchanged in the last summation}) \\
&= \sum_{i_1, \dots, i_{m-1}=1}^n e_{i_1} \otimes \cdots \otimes e_{i_{m-1}} \otimes \\
&\quad [D(\sum_{i=1}^n e_i \otimes T_{i_1}^* \cdots T_{i_{m-2}}^* h_{i_{m-1}i} - \sum_{i,j=1}^n e_j \otimes T_i T_{i_1}^* \cdots T_{i_{m-2}}^* T_j^* h_{i_{m-1}i}) \\
&\quad \{ \sum_{i,j=1}^n e_i \otimes D(e_j \otimes (T_{i_1}^* \cdots T_{i_{m-2}}^* T_j^* h_{i_{m-1}i} - T_{i_1}^* \cdots T_{i_{m-2}}^* T_i^* h_{i_{m-1}j})) \}] \\
&= \sum_{i_1, \dots, i_{m-1}=1}^n e_{i_1} \otimes \cdots \otimes e_{i_{m-1}} \otimes \sum_{i=1}^n D(e_i \otimes T_{i_1}^* \cdots T_{i_{m-2}}^* h_{i_{m-1}i}) \\
&\quad + \sum_{i_1, \dots, i_{m-1}, i, j=1}^n e_{i_1} \otimes \cdots \otimes e_{i_{m-1}} \otimes e_i \otimes \\
&\quad D(e_j \otimes T_{i_1}^* \cdots T_{i_{m-2}}^* (T_j^* T_i^* h_{i_{m-1}} - T_j^* T_{i_{m-1}}^* h_i - T_i^* T_j^* h_{i_{m-1}} + T_i^* T_{i_{m-1}}^* h_j)) \\
&= \sum_{i_1, \dots, i_{m-2}, i, j=1}^n e_{i_1} \otimes \cdots \otimes e_{i_{m-2}} \otimes (e_i \otimes D(e_j \otimes T_{i_1}^* \cdots T_{i_{m-2}}^* h_{ij}) \\
&\quad + \sum_{i_1, \dots, i_{m-1}, i, j=1}^n e_{i_1} \otimes \cdots \otimes e_{i_{m-1}} \otimes e_i \otimes D(e_j \otimes T_{i_1}^* \cdots T_{i_{m-2}}^* (-T_j^* T_{i_{m-1}}^* h_i + T_i^* T_{i_{m-1}}^* h_j)) \\
&\quad (\text{in the term above, index } i_{m-1} \text{ has been replaced by } i \text{ and } i \text{ has been replaced by } j \text{ in the first summation}) \\
&= \sum_{i_1, \dots, i_{m-2}, i, j=1}^n e_{i_1} \otimes \cdots \otimes e_{i_{m-2}} \otimes e_i \otimes D(e_j \otimes T_{i_1}^* \cdots T_{i_{m-2}}^* h_{ij}) \\
&\quad - \sum_{i_1, \dots, i_{m-1}, i, j=1}^n e_{i_1} \otimes \cdots \otimes e_{i_{m-1}} \otimes e_i \otimes D(e_j \otimes T_{i_1}^* \cdots T_{i_{m-1}}^* h_{ij}) = \tilde{x}_{m-1} - \tilde{x}_m.
\end{aligned}$$

So, $\langle y, \tilde{x}_{m-1} - \tilde{x}_m \rangle = 0$.

Next, we would show that $\|\tilde{x}_{m+1}\| = \|\tilde{x}_0\| = 1$ for all $m \in \mathbb{N}$.

$$\begin{aligned}
\|\tilde{x}_{m+1}\|^2 &= \left\langle \sum_{i_1, \dots, i_m, i, j=1}^n e_{i_1} \otimes \dots \otimes e_{i_m} \otimes e_i \otimes D(e_j \otimes T_{i_1}^* \dots T_{i_m}^* h_{ij}), \right. \\
&\quad \left. \sum_{i'_1, \dots, i'_{m-1}, i', j'=1}^n e_{i'_1} \otimes \dots \otimes e_{i'_{m-1}} \otimes e_{i'} \otimes D(e_{j'} \otimes T_{i'_1}^* \dots T_{i'_m}^* h_{i'j'}) \right\rangle \\
&= \sum_{i_1, \dots, i_m, i=1}^n \left\langle \sum_{j=1}^n D(e_j \otimes T_{i_1}^* \dots T_{i_m}^* h_{ij}), \sum_{j'=1}^n D(e_{j'} \otimes T_{i_1}^* \dots T_{i_m}^* h_{ij'}) \right\rangle \\
&= \sum_{i_1, \dots, i_m, i=1}^n \left\langle D\left(\sum_{j=1}^n e_j \otimes T_{i_1}^* \dots T_{i_m}^* h_{ij}\right), \sum_{j'=1}^n e_{j'} \otimes T_{i_1}^* \dots T_{i_m}^* h_{ij'} \right\rangle \\
&= \sum_{i_1, \dots, i_m, i=1}^n \left\langle \sum_{l, k=1}^n e_l \otimes T_k(T_k^* T_{i_1}^* \dots T_{i_m}^* h_{il} - T_l^* T_{i_1}^* \dots T_{i_m}^* h_{ik}), \sum_{j'=1}^n e_{j'} \otimes T_{i_1}^* \dots T_{i_m}^* h_{ij'} \right\rangle \\
&= \sum_{i_1, \dots, i_m, i, j=1}^n \left\langle \sum_{k=1}^n T_k(T_k^* T_{i_1}^* \dots T_{i_m}^* h_{ij} - T_j^* T_{i_1}^* \dots T_{i_m}^* h_{ik}), T_{i_1}^* \dots T_{i_m}^* h_{ij} \right\rangle \\
&= \sum_{i, j=1}^n \left\langle \sum_{k=1}^n (T_k T_k^* h_{ij} - T_k T_j^* h_{ik}), h_{ij} \right\rangle \\
&= \sum_{i, j=1}^n \left\langle \sum_{k=1}^n (T_k T_k^* T_j^* h_i - T_k T_k^* T_i^* h_j - T_k T_j^* T_k^* h_i + T_k T_j^* T_i^* h_k), T_j^* h_i - T_i^* h_j \right\rangle \\
&= \sum_{i, j=1}^n \left\langle \sum_{k=1}^n (T_k T_j^* T_i^* h_k - T_k T_k^* T_i^* h_j), T_j^* h_i - T_i^* h_j \right\rangle \\
&= \sum_{i, j=1}^n \left\langle \sum_{k=1}^n (T_k T_j^* T_i^* h_k) - T_i^* h_j, T_j^* h_i - T_i^* h_j \right\rangle \\
&= \sum_{i, j=1}^n \left\langle T_j \left(\sum_{k=1}^n T_k T_j^* T_i^* h_k \right) - T_j T_i^* h_j, h_i \right\rangle - \sum_{i, j=1}^n \left\langle T_i \left(\sum_{k=1}^n T_k T_j^* T_i^* h_k \right) - T_i T_i^* h_j, h_j \right\rangle \\
&= \sum_{i=1}^n \left\langle \sum_{j=1}^n T_j T_j^* \left(\sum_{k=1}^n T_k T_i^* h_k \right) - \sum_{j=1}^n T_j T_i^* h_j, h_i \right\rangle - \\
&\quad \sum_{j=1}^n \left\langle \sum_{i=1}^n T_i T_i^* \left(\sum_{k=1}^n T_k T_j^* h_k \right) - \sum_{i=1}^n T_i T_i^* h_j, h_j \right\rangle \\
&= \sum_{i=1}^n \left\langle \left(\sum_{k=1}^n T_k T_i^* h_k \right) - \sum_{j=1}^n T_j T_i^* h_j, h_i \right\rangle - \sum_{j=1}^n \left\langle \sum_{k=1}^n T_k T_j^* h_k - h_j, h_j \right\rangle \\
&= \sum_{j=1}^n \left\langle h_j - \sum_{k=1}^n T_k T_j^* h_k, h_j \right\rangle = \langle D(h_1, \dots, h_n), (h_1, \dots, h_n) \rangle = \|\tilde{x}_0\|^2 = 1.
\end{aligned}$$

As $\langle y, \tilde{x}_0 + \tilde{x}_1 \rangle = 0$ and $\langle y, \tilde{x}_m - \tilde{x}_{m+1} \rangle = 0$ for $m \in \mathbb{N}$, we get $\langle y, \tilde{x}_0 + \tilde{x}_{m+1} \rangle = 0$ for $m \in \mathbb{N}$. This implies $1 = \langle \tilde{y}_0, \tilde{y}_0 \rangle = \langle \tilde{y}_0, \tilde{x}_0 \rangle = -\langle \tilde{y}_{m+1}, \tilde{x}_{m+1} \rangle$. By Cauchy-Schwarz inequality, $1 \leq \|\tilde{y}_{m+1}\| \|\tilde{x}_{m+1}\|$, i. e., $1 \leq \|\tilde{y}_{m+1}\|$ for $m \in \mathbb{N}$. This is a contradiction as $y = 0 \oplus \oplus_{m \geq 0} \tilde{y}_m$ is in the Hilbert space $\tilde{\mathcal{H}}$. $\diamond$

4. REPRESENTATIONS OF CUNTZ ALGEBRAS

For $n \geq 2$, the Cuntz algebra $\mathcal{O}_n$ is the C^* -algebra generated by n -isometries $\underline{s} = \{s_1, \dots, s_n\}$, satisfying Cuntz relations: $s_i^* s_j = \delta_{ij} I$, $1 \leq i, j \leq n$, and $\sum s_i s_i^* = I$. It admits many unitarily inequivalent representations. Various classes of representations of $\mathcal{O}_n$ have been constructed in [9, 10], [14]. Given a tuple of contractions $\underline{T} = (T_1, \dots, T_n)$ on a Hilbert space satisfying $\sum T_i T_i^* = I$, we consider its minimal isometric dilation $\underline{\tilde{V}} = (\tilde{V}_1, \dots, \tilde{V}_n)$. We know that the isometries $\tilde{V}_i$ satisfy Cuntz relations and we obtain a representation $\pi_{\underline{T}}$ of the Cuntz algebra $\mathcal{O}_n$ by setting $\pi_{\underline{T}}(s_i) = \tilde{V}_i$. We wish to classify all representations of $\mathcal{O}_n$ we can obtain by dilating *commuting* contractive tuples $\underline{T}$.

Let $\mathcal{S}_n = C(\partial B_n)$ be the C^* -algebra of all continuous complex valued functions on the sphere $\partial B_n = \{(z_1, \dots, z_n) : \sum |z_i|^2 = 1\}$. We have a distinguished tuple $\underline{z} = (z_1, \dots, z_n)$ of elements in $\mathcal{S}_n$ consisting of co-ordinate functions. Given any spherical unitary $\underline{Z} = (Z_1, \dots, Z_n)$ there is a unique representation of $\mathcal{S}_n$ which maps z_i to Z_i . Now given any commuting n -tuple of operators $\underline{T}$, satisfying $\sum T_i T_i^* = I$, we consider its standard commuting dilation $\underline{\tilde{S}} = (\tilde{S}_1, \dots, \tilde{S}_n)$. Let $\rho_{\underline{T}}$ be the representation of $\mathcal{S}_n$, obtained by taking $\rho_{\underline{T}}(z_i) = \tilde{S}_i$.

DEFINITION 16. Let π be representation of $\mathcal{O}_n$ on a Hilbert space $\mathcal{L}$ with $\underline{W} = (W_1, \dots, W_n) = (\pi(s_1), \dots, \pi(s_n))$. The representation π is said to be *spherical* if $\overline{\text{span}} \{ \underline{W}^\alpha h : h \in \mathcal{L}^c(\underline{W}), \alpha \in \Lambda \} = \mathcal{L}$, where $\mathcal{L}^c(\underline{W})$ is the space where the maximal commuting piece $\underline{W}^c$ of $\underline{W}$ acts as in Definition 3.

Note that this Definition means in particular that if π is spherical then the maximal commuting piece $\underline{W}^c$ is non-trivial. We will see that it is actually a spherical unitary. But this is not a justification for calling such representations as spherical, because this happens for any representation of $\mathcal{O}_n$, as long as $\underline{W}^c$ is non-trivial! The actual justification of this Definition is in Theorem 18.

THEOREM 17. Let $\underline{T} = (T_1, \dots, T_n)$ be a commuting tuple of operators on a Hilbert space $\mathcal{H}$, satisfying $\sum T_i T_i^* = I$. Then the representation $\pi_{\underline{T}}$ coming from the minimal isometric dilation of $\underline{T}$ is spherical. Suppose $\underline{R} = (R_1, \dots, R_n)$ is another commuting tuple, possibly on a different Hilbert space, satisfying $\sum R_i R_i^* = I$. Then the representations $\pi_{\underline{T}}$, $\pi_{\underline{R}}$ of $\mathcal{O}_n$ are unitarily equivalent if and only if the representations $\rho_{\underline{T}}$, $\rho_{\underline{R}}$ of $\mathcal{S}_n$ are unitarily equivalent.

PROOF: In view of Theorem 13, the maximal commuting piece of the minimal isometric dilation $\underline{\tilde{V}}$ of $\underline{T}$ is a realization of the standard commuting dilation $\underline{\tilde{S}}$ of $\underline{T}$. The first claim follows easily as the space on which the standard commuting dilation acts includes the original space $\mathcal{H}$. So $\underline{\tilde{V}}$ is the minimal isometric dilation of $\underline{\tilde{S}}$. Similar statement holds for the tuple $\underline{R}$. Now the Theorem follows due to uniqueness up to equivalence of minimal isometric dilation of contractive tuples, and unitary equivalence of maximal commuting pieces of unitarily equivalent tuples. $\diamond$

So this Theorem reduces the classification problem for representations of $\mathcal{O}_n$ arising out of general commuting tuples to that of representations of $\mathcal{S}_n$. But $\mathcal{S}_n$ being a commutative C^* -algebra, its representations are well-understood and is part of standard C^* -algebra theory. We find the description of this theory as presented in Arveson's classic [4] most suitable for our purposes.

Given any point $w = (w_1, \dots, w_n) \in \partial B_n$, we have a one dimensional representation ϕ_w of $\mathcal{S}_n$, which maps f to $f(w)$. Of course w is a spherical unitary as operator tuple on $\mathbb{C}$. We can construct the minimal isometric dilation $(W_1^w, \dots, W_n^w)$ of this tuple as in the proof of Theorem 15 (Schäffer construction). We see that the dilation space is

$$\mathcal{H}^w = \mathbb{C} \oplus (\Gamma(\mathbb{C}^n) \otimes \mathbb{C}_w^n) \subseteq \mathbb{C} \oplus (\Gamma(\mathbb{C}^n) \otimes \mathbb{C}^n),$$

where $\mathbb{C}_w^n$ is the subspace of vectors orthogonal to $(\overline{w_1}, \dots, \overline{w_n})$ in $\mathbb{C}^n$. Further the operators W_i^w are given by

$$W_i^w(h \oplus \sum_{\alpha} e^{\alpha} \otimes d_{\alpha}) = w_i h \oplus D(e_i \otimes h) \oplus e_i \otimes (\sum_{\alpha} e^{\alpha} \otimes d_{\alpha}).$$

We denote the associated representation of $\mathcal{O}_n$ by ρ_w . This representation is known to be irreducible as it is nothing but the GNS representation of the so-called Cuntz state on $\mathcal{O}_n$ (See [14], Example 5.1)], given by

$$s_{i_1} \cdots s_{i_m} s_{j_1}^* \cdots s_{j_p}^* \mapsto w_{i_1} \cdots w_{i_m} \overline{w_{j_1}} \cdots \overline{w_{j_p}}.$$

Now an arbitrary multiplicity free representation of $\mathcal{S}_n$ can be described as follows [4]. Consider a finite Borel measure μ on ∂B_n . Then we get a representation of $\mathcal{S}_n$ on the Hilbert space $L^2(\partial B_n, \mu)$, which sends $f \in \mathcal{S}_n$ to the operator 'multiplication by f '. This representation can be thought of as direct integral of representations ϕ_w with respect to measure μ . Now it is not hard to see that the associated representation of $\mathcal{O}_n$ is simply the direct integral of representations ρ_w with respect to measure μ and acts on $\oint \mathcal{H}^w \mu(dw)$. Finally an arbitrary representation of $\mathcal{S}_n$ is a countable direct sum of such multiplicity free representations. So we have proved the following result.

THEOREM 18. *Every spherical representation of $\mathcal{O}_n$ is a direct integral of representations $\rho_w, w \in \partial B_n$ (GNS representations of Cuntz states).*

Here we have not bothered to write down as to when two such representations are equivalent. But in view of Theorem 17, we can do it exactly as in ([4], page:54-55), by keeping track of multiplicities and equivalence classes of measures.

THEOREM 19. *Let π be a representation of $\mathcal{O}_n$. Then (i) π decomposes uniquely as $\pi = \pi^0 \oplus \pi^1$, where π^0 is spherical and $(\pi^1(s_1), \dots, \pi^1(s_n))$ has trivial maximal commuting piece (Either π^0 or π^1 could also be absent); (ii) The maximal commuting piece of $(\pi(s_1), \dots, \pi(s_n))$ is either trivial or it is a spherical unitary. (iii) If π is irreducible then either the maximal commuting piece is trivial or it is one dimensional. In the second case, it is unitarily equivalent to GNS representation of a Cuntz state.*

PROOF: Suppose π is a representation of $\mathcal{O}_n$ on a Hilbert space $\mathcal{L}$ and $\underline{W} = (\pi(s_1), \dots, \pi(s_n))$. Consider the space $\mathcal{L}^0$ generated by $\mathcal{L}^c(\underline{W})$ as $\mathcal{L}^0 = \overline{\text{span}} \{ \underline{W}^{\alpha} h : h \in \mathcal{L}^c(\underline{W}), \alpha \in \tilde{\Lambda} \}$. Now each W_i^* leaves $\mathcal{L}^c(\underline{W})$ invariant and clearly $\mathcal{O}_n = C^* \{ \underline{s}^{\alpha} (\underline{s}^{\beta})^* : \alpha, \beta \in \tilde{\Lambda} \}$. Then it follows that $\mathcal{L}^0$ is a reducing subspace for π . Taking $\mathcal{L}^1 = (\mathcal{L}^0)^{\perp}$, we decompose π as $\pi^0 \oplus \pi^1$ with respect to $\mathcal{L} = \mathcal{L}^0 \oplus \mathcal{L}^1$. It is clear that this is a decomposition as required by (i). Uniqueness of this

decomposition and (ii) follow easily as maximal commuting piece of direct sum of tuples is direct sum of maximal commuting pieces (Corollary 5) and then (iii) follows from Theorem 18. $\diamond$

Let us see as to what happens if we dilate commuting tuples $\underline{T}$, satisfying just $\sum T_i T_i^* \leq I$. In this case, as is well-known, the minimal isometric dilation decomposes as $((V_1 \otimes I) \oplus W_1, \dots, (V_n \otimes I) \oplus W_n)$ where $(V_1, \dots, V_n)$ is the standard tuple of full Fock space, and $(W_1, \dots, W_n)$ are isometries satisfying Cuntz relations. If $\underline{T}$ is not pure the term $(W_1, \dots, W_n)$ is present and we get a representation of $\mathcal{O}_n$. However, as seen in the proof of Theorem 13, $(W_1, \dots, W_n)$ is a minimal isometric dilation of a spherical tuple $(Z_1, \dots, Z_n)$ (the ‘spherical part’ of the standard commuting dilation of $\underline{T}$) and hence the representation of $\mathcal{O}_n$ we get is still spherical.

Finally we remark that it is easy to get examples of non-commuting tuples dilating to representations of $\mathcal{O}_n$ which are not spherical. For instance we can consider the tuple $\underline{R} = (R_1, R_2)$ on $\mathbb{C}^2$ defined by

$$R_1 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, R_2 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}.$$

Then as $R_1 R_1^* + R_2 R_2^* = 1$, the minimal isometric dilation of (R_1, R_2) satisfies Cuntz relations. We can see that it has trivial commuting piece through a simple application of Corollary 4.3 of [14].

Acknowledgements: The first author is supported by Indo-French (IFCPAR) grant No: IFC/2301-1/2001/1253, the second author is supported by research grant no. SR/FTP/MS-16/2001 of the Department of Science and Technology, India and the last author is supported by a research fellowship from the Indian Statistical Institute.

REFERENCES

- [1] J. AGLER, ‘The Arveson extension theorem and coanalytic models’, *Integral Equations and Operator Theory*, 5 (1982) 608-631. **MR 84g:47011.**
- [2] A. ARIAS and G. POPESCU, ‘Noncommutative interpolation and Poisson transforms,’ *Israel J. Math.*, 115 (2000) 205-234. **MR 2001i:47021.**
- [3] A. ARIAS and G. POPESCU, ‘Noncommutative interpolation and Poisson transforms II’, *Houston J. Math.*, 25 (1999) 79-98. **MR 2000i:47148.**
- [4] W. B. ARVESON, *An Invitation to C^* -algebras*, Graduate Texts in Mathematics, No. 39, (Springer-Verlag, New York-Heidelberg, 1976). **MR 58#23621.**
- [5] W. B. ARVESON, ‘Subalgebras of C^* -algebras III, Multivariable operator theory,’ *Acta Math.*, (2) 181 (1998) 159-228. **MR 2000e:47013.**
- [6] A. ATHAVALA, ‘On the intertwining of joint isometries,’ *J. Operator Theory*, 23 (1990) 339-350. **MR 91i:47029.**
- [7] A. ATHAVALA, ‘Model theory on the unit ball in $\mathbb{C}^m$,’ *J. Operator Theory*, 27 (1992) 347-358. **MR 94i:47011**
- [8] B. V. RAJARAMA BHAT and T. BHATTACHARYYA, ‘A model theory for q -commuting contractive tuples’, to appear in the *J. Operator Theory* (2002).
- [9] O. BRATTELI and PALLE E. T. JORGENSEN *Iterated function systems and permutation representations of the Cuntz algebra*, Mem. Amer. Math. Soc., (663) 139 (1999). **MR 99k:46094a.**

- [10] O. BRATTELI and PALLE E. T. JORGENSEN, 'Isometries, shifts, Cuntz algebras and multiresolution wavelet analysis of scale N ', *Integral Equations Operator Theory*, 28 (1997) 382-443. **MR 99k:46094b**.
- [11] J. W. BUNCE, 'Models for n -tuples of noncommuting operators,' *J. Funct. Anal.*, 57 (1984) 21-30. **MR 85k:47019**
- [12] J. CUNTZ, 'Simple C^* -algebras generated by isometries,' *Commun. Math. Phys.*, 57 (1977) 173-185. **MR 57#7189**.
- [13] C. DAVIS, 'Some dilation and representation theorems', *Proceedings of the Second International Symposium in West Africa on Functional Analysis and its Applications* (Kumasi, 1979), pp.159-182. **MR 84e:47012**.
- [14] K. R. DAVIDSON, D. W. KRIBS, and M. E. SHPIGEL, 'Isometric dilations of noncommuting finite rank n -tuples', *Canad. J. Math.*, 53 (2001) 506-545.
- [15] S. W. DRURY, 'A generalization of von Neumann's inequality to the complex ball', *Proc. Amer. Math. Soc.*, 68 (1978) 300-304. **MR 80c:47010**.
- [16] A. E. FRAZHO, 'Models for noncommuting operators', *J. Funct. Anal.*, 48 (1982) 1-11. **MR 84h:47010**.
- [17] A. E. FRAZHO, 'Complements to models for noncommuting operators', *J. Funct. Anal.*, 59 (1984) 445-461. **MR 86h:47010**.
- [18] P. R. HALMOS, *A Hilbert Space Problem Book*, Second Edition, Graduate Texts in Mathematics No. 19, (Springer-Verlag, New York-Berlin 1982). **MR 84e:47001**.
- [19] S. PARROTT, 'Unitary dilations for commuting contractions', *Pacific J. Math.*, 34 (1970) 481-490. **MR 42#3607**.
- [20] G. POPESCU, 'Isometric dilations for infinite sequences of noncommuting operators', *Trans. Amer. Math. Soc.*, 316 (1989) 523-536. **MR 90c:47006**
- [21] G. POPESCU, 'Models for infinite sequences of noncommuting operators', *Acta Sci. Math. (Szeged)*, 53 (1989) 355-368. **MR 91b:47025**.
- [22] G. POPESCU, 'Characteristic functions for infinite sequences of noncommuting operators', *J. Operator Theory*, 22 (1989) 51-71. **MR 91m:47012**.
- [23] G. POPESCU, 'Poisson transforms on some C^* -algebras generated by isometries', *J. Funct. Anal.*, 161 (1999) 27-61. **MR 2000m:46117**.
- [24] G. POPESCU, 'Curvature invariant for Hilbert modules over free semigroup algebras', *Advances in Mathematics*, 158 (2001) 264-309. **MR 2002b:46097**.
- [25] C. R. PUTNAM, *Commutation properties of Hilbert space Operators and Related Topics*, (Springer-Verlag, New York, 1967). **MR 36#707**.
- [26] J. J. SCHÄFFER, 'On unitary dilations of contractions', *Proc. Amer. Math. Soc.*, 6 (1955) 322. **MR 16,934c**

B. V. RAJARAMA BHAT AND SANTANU DEY
 Indian Statistical Institute,
 R. V. College Post,
 Bangalore 560059, India.
 e-mail: bhat@isibang.ac.in
 and santanu@isibang.ac.in

TIRTHANKAR BHATTACHARYYA
 Department of Mathematics,
 Indian Institute of Science
 Bangalore 560012, India.
tirtha@math.iisc.ernet.in